Everwipe™ Disinfectant Wipes will clean and disinfect3 hard, non-porous surfaces of cabinets, chairs, computers, countertops, desks, desktops, doorknobs, carts, garbage cans, headsets, keyboards, railings, seats, sinks, tables, tabletops, telephones, trashcans, exercise machines, faucets, bathtubs, shower doors, light switches, DIRECTIONS FOR LISE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling This product is not for use on critical and semi-critical medical device surfaces.

CLEANING: Use one towelette to completely remove all dirt and debris from surfaces to be treated

DISINFECTING: To disinfect hard, non-porous surfaces, thoroughly wet the surface to be treated. Treated surface must remain visibly wet for disinfecting. Use enough towelettes for treated surface to remain visibly wet for 5 minutes. For visibly soiled surfaces, clean before following disinfecting instructions. Allow surfaces to air dry. This product is an effective disinfectant against Escherichia coli (E. coli) O157:H7. Pseudomonas aeruginosa. Salmonella enterica (Salmonella), Staphylococcus aureus, Streptococcus pyogenes (Strep), Vancomycin resistant Enterococcus faecium (VRE), Methicillin resistant Staphylococcus aureus (MRSA), Community Associated Methicillin resistant Staphylococcus aureus (CA-MRSA) Genotype USA400, Community Associated Methicillin resistant Staphylococcus aureus (CA-MRSA) Genotype USA300, Vancomycin intermediate resistant Staphylococcus aureus (VISA) and Klebsiella pneumoniae New Dehli Metallo-Beta Lactamase (NDM-1) Carbapenem Resistant (CDC Clinical Isolate No. 10002).

#### EFFICACY TESTS HAVE DEMONSTRATED THAT THIS PRODUCT IS AN EFFECTIVE BACTERICIDE, VIRUCIDE AND FUNGICIDE IN THE PRESENCE OF ORGANIC SOIL (5% BLOOD SERUM).

SANITIZING/DEODORIZING: To sanitize/deodorize hard, non-porous surfaces; wipe surface, using enough towelettes to allow treated surface to remain wet for 15 seconds for Staphylococcus aureus (ATCC 6538), Klebsiella pneumoniae (ATCC 4352), Listeria monocytogenes (ATCC 35152), Escherichia coli O157:H7 (ATCC 43895) and 60 seconds for Campylobacter jejuni (ATTC 29428). Let surface dry. For visibly soiled surfaces, clean excess dirt first.

For surfaces that may come in contact with food, a potable water rinse is required. For plastic and painted surfaces, spot test on an inconspicuous area before use. This product will not harm most surfaces including acrylic, fiberglass and vinyl. <u>Do not use on upainted</u> wood. Not for personal cleansing. This is <u>not</u> a baby wipe! Do not flush in toilet.

\*VIRUCIDAL ACTIVITY: This product is an effective virucide on hard, non-porous surfaces against Bovine Viral Diarrhea Virus (BVDV), Influenza A2/Japan/305/57 Virus (H2N2), Hepatitis B Virus (HBV), Hepatitis C Virus (HCV), Duck Hepatitis B Virus (DHBV), pes Simplex Virus 1 (HSV-1) (ATCC VR-260), Herpes Simplex Virus 2 (HSV-2) (ATCC VR-734), Human Immunodeficiency Virus Type 1 (HIV-1) when the treated surface is allowed to remain wet for **5 minutes**, Human Coronavirus (ATTC VR-740, strain 229E) when the treated surface is allowed to remain wet for 10 minutes, and SARS-Related Coronavirus 2 ISARS-CoV-21 USA-WA1/2020 BEI NR-52281 when the treated surface is allowed to remain wet for 30 seconds

FUNGICIDAL ACTIVITY: This product is an effective fungicide against Trichophyton interdigitale, formerly Trichophyton mentagrophytes (ATCC 9533), causative agent of Athlete's Foot Fungus, when used as directed on hard surfaces found in bathrooms, shower stalls, locker rooms, exercise facilities or other clean, hard, non-porous, surfaces commonly contacted by bare feet. The surface must remain visibly wet for 3 minutes. Allow surfaces to air dry.

Manufactured by: Legacy Converting, Inc. 1 Advantage Court, Bordentown, NJ 08505 www.everwipe.com 800.521.4190

## <u>.egacy</u>

EPA Reg. No. 1839-190-88331 EPA Est. No. 88331-NJ-1

# **EVERWIPE**"

### DISINFECTANT WIPES

## Kills SARS-CoV-2 (COVID-19 virus)

on hard non-porous surfaces in 30 seconds!

#### KILLS 99.9% OF BACTERIA IN JUST 15 SECONDS!\*\* | KILLS COLD' & FLU VIRUSES\*\*\*

## **Cleans, Disinfects<sup>3</sup> and Deodorizes**

..<u>99.72%</u>

#### NGREDIENTS % C14, 30% C16, 5% C12, 5% C18) C12. 32% C14) m chlorides.....0.14% ethylhenzyl amm IGREDIENTS

PRECAUTIONARY STATEMENTS.

NET CONTENTS: 1.18lbs (0.535kg) | 7" x 7" (18cm x 18cm)

Kills Flu Virustt, Kills HIV-1, Kills Antibiotic Resistant Bacteria: CA-MRSA, MRSA, VISA, VRE and NDM-1 Kills Campylobacter jejuni in just 60 seconds! Disinfects <sup>#</sup>household, <sup>#</sup>kitchen and/or bathroom bacteria<sup>†</sup>

\* Staphylococcus aureus (Staph), Klebsiella pneumoniae, Listeria monocytogenes, Escherichia coli (E. coli)

75

<sup>1</sup> Pseudomonas aeruginosa, Staphylococcus aureus (Staph), Salmonella enterica (Salmonella) Streptococcus pyogenes (Strep), Escherichia coli (E. coli) O157:H7 †† Influenza Ao*i*, lánan

Influenza A Virus, Pandemic 2009 H1N1 Influenza A Virus, Human Coronavirus<sup>1</sup>

Human Coronavirus (ATCC VR-740) #When using on food contact surfaces: thoroughly rinse all treated surfaces with potable (tap) water. <sup>3</sup> When used according to disinfection directions

#### KILLS HIV-1 ON PRE-CLEANED ENVIRONMENTAL SURFACES/ **OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS**

in health care settings (Hospitals, Nursing Homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1, Human Hepatitis B Virus (HBV) and Human Hepatitis C Virus (HCV).

#### SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV and HCV OF SURFACES / OBJECTS SOILED WITH BLOOD/BODY FLUIDS

PERSONAL PROTECTION: When handling items soiled with blood or body fluids, use disposable latex gloves, gowns, masks and eye coverings. CLEANING PROCEDURES: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product. CONTACT TIME: Allow surface to remain visibly wet for 5 minutes. DISPOSAL OF INFECTIOUS MATERIALS: Blood and other body fluids should be autoclaved and disposed of according to local regulations for infectious waste disposal.

#### TO OPEN PACKAGE

1. Lift tab at front of lid to open. 2. Pull up corner of center sheet and thread through dispenser slit in lid. 3. Pull out exposed towelette at a 45 degree angle and snap off. The next towelette is ready for dispensing. 4. Close lid to retain moisture. 5. Dispose of used towelettes in trash. Do not flush.

#### STORAGE AND DISPOSAL:

Do not contaminate water, food or feed by storage or disposal CONTAINER STORAGE/DISPOSAL: Store in an area inaccessible to children. Wipes: Dispose of towelette in trash after use. Do not flush. Canister: Tightly close lid between uses to retain moisture. Nonrefillable container. Do not reuse or refill this container. Offer empty container for recycling if available or discard in trash.

#### PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling

FIRST AID.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

IF IN EYES: Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.





#### item# 101075



KEEP OUT OF REACH OF CHILDREN CAUTION	ACTIVE II Alkyl (60% dimethyl b Alkyl (68% dimethyl e INERT IN
SEE BACK PANELS FOR ADDITIONAL	TOTAL